

Toward the dynamic interactome: it's about time

Teresa M. Przytycka, Mona Singh and Donna K. Slonim

Submitted: 20th July 2009; Received (in revised form): 1st November 2009

Abstract

Dynamic molecular interactions play a central role in regulating the functioning of cells and organisms. The availability of experimentally determined large-scale cellular networks, along with other high-throughput experimental data sets that provide snapshots of biological systems at different times and conditions, is increasingly helpful in elucidating interaction dynamics. Here we review the beginnings of a new subfield within computational biology, one focused on the global inference and analysis of the dynamic interactome. This burgeoning research area, which entails a shift from static to dynamic network analysis, promises to be a major step forward in our ability to model and reason about cellular function and behavior.

Keywords: network analysis; network dynamics; interaction networks; systems biology

INTRODUCTION

Over the past decade, high-throughput experimental and computational methods have been developed to infer and predict the structure of gene and protein networks. As a result, large-scale cellular networks have been obtained for a wide range of organisms across the evolutionary spectrum [1–12]. Computational analyses of these networks have great potential in aiding our understanding of gene function, biological pathways and cellular organization. While significant progress has been made in computational analysis of proteome-scale cellular networks [13, 14], the dynamics inherent within these networks are often overlooked in computational network analysis. However, proper cellular functioning requires the precise coordination of a large number of events, and identifying the temporal and contextual signals underlying proposed interactions is a crucial part of understanding cellular function. Network dynamics can describe, for example, how cells respond to environmental cues or how an interaction network changes during development or differentiation.

We hold the view that modeling and analyzing interaction and network dynamics should be at the forefront of current research efforts in computational interactomics.

The first question, perhaps, is to define what we mean by interaction or network ‘dynamics’. In a simplified view, an interaction may occur or not depending upon spatial, temporal and/or contextual variation. Though temporal variation of interactions can be assessed on an evolutionary time scale, in this review we focus primarily on short-term interaction variation within a specific organism. Interactions may be *constitutive* or *obligate*, or may instead occur only in specific situations. Among these dynamically varying interactions (sometimes referred to as *transient* interactions), the variation may be either *reactive* (i.e. caused by exogenous factors, such as a response to some environmental stimulus) or *programmed* (i.e. due to endogenous signals, such as cell-cycle dynamics or developmental processes). Contextual variation overlaps heavily with temporal variation, but focuses more specifically on characterizing

Corresponding author. Teresa M. Przytycka, National Center of Biotechnology Information, NLM, NIH, 8000 Rockville Pike, Bethesda MD 20814, USA. Tel: (301) 402-172; Fax: (301) 480-4837; E-mail: przytyck@ncbi.nlm.nih.gov

Teresa Przytycka is an Investigator at the National Center of Biotechnology Information, NLM, NIH, where she heads the research group ‘Algorithms and Graph Theory for Computational and Systems Biology’.

Donna Slonim is an Associate Professor in the Department of Computer Science at Tufts University and in the Department of Pathology at Tufts University School of Medicine.

Mona Singh is an Associate Professor at Princeton University in the Department of Computer Science and the Lewis-Sigler Institute for Integrative Genomics.

All authors contributed equally to this work.

reactive variation and the conditions that cause it. Studying context may also encompass examining sequence or genetic variation within a population of contemporaries and exploring how that variation affects network interactions, topology and function.

Of course, biologists have studied dynamics in biological systems for many years. Traditionally, efforts have focused on individual genes or proteins as well as specific interactions in limited contexts. At a somewhat larger scale, metabolic networks, which inherently include temporal information and rely on the availability of specific metabolites as context, have also been mapped across numerous organisms [15]. However, within the last few years, we have been given an unprecedented opportunity to investigate molecular networks from a global perspective, using diverse high-throughput experimental data to inform our understanding of molecular interactions on a genomic scale. Our goal in this review is to focus specifically on research that links these global analyses to the dynamics and context of biological systems. We will highlight bioinformatics approaches that utilize genomic-scale experimental data sets for analyzing network dynamics rather than more established methods for quantitative modeling and simulation of dynamical systems [16–18] or their more recent extensions and implementations [19–26].

If dynamics play such a central role in biological systems, why have genome-scale computational analyses of interaction and network dynamics remained elusive? One central reason is that the most widely-applied large-scale technologies to determine protein interactions, such as yeast two-hybrid [27] and TAP-MS [28] to detect protein–protein interactions or *in vitro* proteome microarrays to detect phosphorylation interactions [12], do not provide spatial, temporal or contextual information about detected interactions. ChIP–chip [29] or ChIP–seq [30] approaches for uncovering regulatory interactions have been used to uncover reactive contextual variation [10] but have only recently started to be used to uncover temporal variation over a dynamic time course [31, 32]. Thus, even if all possible interactions in an organism could be determined using these technologies, for any given protein, it would generally not be known when and where each of its interactions occurs.

Computational approaches provide one means for inferring and analyzing changes in interaction and network dynamics. Broadly speaking, dynamically

changing interactions can be thought of as a means for transmitting ‘information’ through cellular networks, and much of the work we discuss here attempts to characterize this ‘flow’ of information, often without an explicit understanding of the full dynamic nature of individual interactions. Specific research goals toward computationally elucidating interaction and network dynamics are diverse. To date, these have included discovering dynamic pathway information from network data, inferring network structure utilizing dynamic data, discerning large-scale changes in network topology and function, and understanding network responses to engineered perturbations (such as knock-outs and knock-downs) or to evolutionary perturbations (such as sequence variation between individuals). We have organized our review with respect to this seemingly broad set of research goals. Ultimately, however, all of the research described here is intended to improve our understanding of biological and physiological processes at the molecular level by attempting to characterize and analyze the dynamic interactome.

PATHWAY INFERENCE FROM PROTEIN INTERACTION NETWORKS

By itself, a complete cellular interaction map gives no information about the ordering of proteins with respect to their regulatory or physical relationships. One way of inferring dynamics from static network data is to use prior biological knowledge to help order the interactions. With sufficient information, one can construct a pattern representing some important biological structure and then search for instances of that pattern in static networks. Signaling pathways provide excellent opportunities for such approaches.

Signaling pathways transfer information detected at the cell surface to regulatory factors in the nucleus in order to orchestrate a cellular response. Steffen *et al.* [33] introduced a computational approach for discovering signaling pathways from protein–protein interaction data. Their approach is based on enumerating relatively short linear paths starting at membrane proteins and ending with DNA-binding proteins. These pathways are then evaluated with respect to gene expression data, with the expectation that proteins in the same pathway should be expressed in the same conditions and at

approximately the same time. Interaction reliabilities can also be incorporated in this approach, along with improved algorithms for path enumeration [34]. While it is convenient to think of one signaling pathway at a time, numerous signals are detected essentially simultaneously by various sensory proteins at the cell surface. Supper *et al.* [35] suggest an approach for inferring signaling pathways based on an arbitrary number of sensor and regulatory proteins, and use Steiner tree formulations that favor bow tie architectures [36] corresponding to intermediate ‘integrator’ core proteins.

Signaling and regulatory pathways, as well as other complex network patterns, may also be described with respect to the properties of their constituent proteins—for example, kinases, transcription factors or descriptions of constituent interaction domains. Banks *et al.* [37] introduced *network schemas* to describe such patterns and developed fast algorithms for searching for their matches in interactomes. A network schema consists of descriptions of proteins (e.g. their molecular functions or putative domains) along with the desired topology and types of interactions (e.g. physical, phosphorylation or regulatory). Network schemas can thus describe the linear patterns underlying signaling pathways suggested in Steffen *et al.* as well as more complex pathways. In addition to searching for matches to particular network schemas, it is also possible to infer which network schemas are frequent and over-represented in networks [38] and to thereby uncover general recurring patterns underlying a range of biological processes.

While the above-mentioned approaches consider signaling pathways in terms of orderings of individual molecules, Zotenko *et al.* [39] proposed an approach that focuses on ordering overlapping groups of molecules, putatively corresponding to transient complexes or functional groups. They approximate signaling networks by chordal graphs [40] in which functional groups, corresponding to dense subgraphs in the original network, are represented as cliques. Then they use clique tree representations of chordal graphs [41] to elucidate (partial) orderings within these functional groups.

Comparative interactomics is an alternative approach for discovering pathways within cellular networks. The core idea is that if a pathway is known in one organism, searching for homologs of its component proteins, along with conserved patterns of interactions, is a powerful technique for

pathway detection [42]. Much research in recent years has focused on such pathway querying as well as the more general task of uncovering conserved pathways and modules via network alignment [42–47].

PATHWAY INFERENCE BY INTEGRATING DYNAMIC DATA WITH PROTEIN INTERACTION NETWORKS

Protein–protein physical and regulatory interaction networks provide a ‘scaffold’ by which to interpret other types of dynamic data and thereby obtain hints about signaling and regulatory pathways. Since proteins along a biological pathway tend to be co-expressed, one approach for identifying pathways is to identify *active subnetworks*, connected regions in physical interaction networks that exhibit significant expression changes over some set of conditions [48]. These attempts to characterize contextual variation in networks have been further extended and improved [49–51]. Such context-specific subnetworks do not have temporal orderings between proteins; combining these subnetworks with the pathway-ordering approaches (such as those reviewed above) may be a promising approach for refining the flow of information through the network.

Expression data can also be utilized to infer causality as well as information flow within cellular networks. A particularly illuminating source of dynamic data comes from knock-out experiments, where a gene is perturbed or removed from a genetic background and the expression levels of all other genes are measured. Yeang *et al.* [52] develop a probabilistic approach for explaining observed gene expression changes due to a knock-out by inferring molecular cascades of ‘flow’ through the interaction network. These molecular cascades correspond to paths beginning from the knock-out gene (‘cause’) and ending at the gene whose expression has changed (‘effect’). Additionally, interactions along the path are inferred to be either activating or repressing, and together must be consistent with the sign of the knock-out effect. The basic approach was later refined and experimentally validated in yeast [53]. Furthermore, it is possible to change the formulation of the problem so that it favors explaining many cause–effect pairs by few pathways [54].

RNA interference (RNAi) screens are a powerful technique for obtaining perturbation data in higher

organisms. Here, a known gene from a pathway of interest is chosen as a reporter gene, other genes in the genome are systematically knocked-down using RNAi, and the effect on the reporter is measured. Flow-based methods through protein interaction networks have been utilized to connect and order genes that affect the reporter [55]. Alternatively, genes that affect the reporter gene, or some other experimentally measurable phenotype of interest, can be used as ‘seeds’ within known protein interaction networks to uncover subnetworks related to the biological process of interest [56]. Methodologically, this approach is similar to the active subnetwork approaches outlined above. As RNA-based loss of function screens are increasingly being applied with automated image analysis to detect effects on specific processes or phenotypes [57, 58], these types of network analyses are likely to be relevant to study a broad range of interesting biological questions. Future challenges for all these types of approaches include incorporating additional types of data into these models (e.g. known annotations of individual proteins or other types of interaction data).

INFERRING NETWORK STRUCTURE FROM DYNAMIC DATA

The earliest attempts to incorporate dynamic data into models of gene or protein interaction networks were aimed primarily at inferring the network structure itself. (See [48, 59–63] for a general introduction to this area.) Most of these early network inference papers do not model dynamics. However, many used dynamic data, most commonly expression data from genome-wide microarrays, to infer static network structure. Such work includes efforts at reconstructing regulatory networks using mutual-information measures [64, 65] or Bayesian/graphical modeling methods [61, 66, 67]. Though reverse-engineering efforts have more frequently focused on inferring regulatory networks, Bayesian networks have been used to infer human T-cell signaling networks from single-cell measurements of phosphorylation in response to perturbations [68]. Parameterized biochemical modeling of responses, as measured by microarrays, to RNA interference and over-expression screens have also been utilized to infer signaling pathways in *Drosophila* [69].

Friedman *et al.* [70] and Murphy and Mian [71] were among the first to suggest using Dynamic Bayesian Networks (DBNs) for modeling regulatory

networks because such models can capture time-dependent structures (such as feedback loops) impossible to express with traditional probabilistic networks. Ong [72] and Kim [73] also explored DBN approaches. However, the aim was typically to infer a static network structure from dynamic data, rather than using the dynamic model to assess the context of specific interactions.

Coexpression data can also be used to construct networks relating genes by the similarity of their expression patterns under different conditions [74, 75]. Coexpression networks have been applied to finding functional modules in expression data [75, 76]. While typically intended to reveal protein function, such networks are directly related to the context of the samples used in their construction. Thus, this information can be exploited explicitly to reveal context-specific patterns in expression data. Choi *et al.* [77] constructed two distinct coexpression networks using data sets representing different conditions (tumors and normal control samples) and identified context-specific differences by comparing the resulting ‘static’ networks. Kostka and Spang [78] proposed an approach for finding sets of genes whose coexpression patterns differed in different contexts. Fang *et al.* [79] have extended this idea to identify ‘subspace’ patterns of differential coexpression (subsets of samples and genes where the genes are coexpressed in most samples from one class but not coexpressed in most samples from the other class). Recent work has also used dynamic modeling to identify transient coexpression relationships capturing static but temporally related snapshots of the dynamic network [80, 81].

By exploiting the modular nature of cellular function, it is possible to infer context-specific regulators from expression data. Groups of genes that are coexpressed in a subset of a conditions are first identified by biclustering approaches [82, 83], and then their condition-specific regulators can be inferred by either relating the expression levels of the genes comprising the module to those of its regulators [84, 85] or by integrating computational or experimental information about regulatory interactions [86]. De Bivort *et al.* take a coarser-grain view of regulatory networks by considering regulatory relationships between the modules themselves and relating expression levels. For one module to expression levels of another module after a time delay [87].

Integration of additional data types has improved the power of network inference methods [88].

Integration of static network and dynamic experimental data was essential for the work of Bromberg *et al.* [89]. Using a mouse network assembled from multiple sources, they predicted a role for the breast cancer protein BRCA1 in neuronal differentiation and suggested a new signaling pathway. Furthermore, by combining this data with transcription factor activation experiments, they have been able to propose a Boolean logic-type description of the neurite outgrowth network. Similarly, Baugh *et al.* [90] integrated temporal and spatial variation to identify a developmental regulatory network in a *Caenorhabditis elegans* model.

The integration of expression data and ChIP–chip data is one particularly promising approach for elucidating dynamic control in regulatory networks. Recently Seok *et al.* used Network Component Analysis (NCA) [91] to build a series of time-varying transcriptional networks for lipopolysaccharide response in humans [92]. NCA is a computational approach that can be used to predict dynamic transcription factor activity over time. It utilizes (partial) knowledge of a regulatory network as well as gene expression data, and its goal is to infer both the concentrations of active transcription factors as well as their promoter affinities. The dynamic networks built by Seok *et al.* captured changes in transcription factor activities and gene expression levels, as well as in signaling and regulatory interactions. Ye *et al.* [93] further extended NCA by developing a computational approach to model the impact of single nucleotide polymorphisms (SNPs) on the concentrations of transcription factors and their promoter affinities. This analysis helps explain how genetic variations linked to gene expression via eQTL analysis (described later in this review) might perturb the regulatory network.

An interesting approach for integrating expression and ChIP–chip data was taken by Ernst *et al.* [94]. They exploited time–course expression experiments to build a detailed dynamic regulatory map for yeast stress response by incorporating a hidden Markov model to identify dynamic ‘bifurcation points’ where expression of a subset of genes diverges from that of others. Recent work by Ucar *et al.* [95] further integrates context-specific ChIP–chip and expression data with context-invariant sequence and nucleosome occupancy data to predict which binding relationships are functional and to characterize the functional roles played by individual transcription factors.

UNCOVERING ORGANIZATIONAL PRINCIPLES UNDERLYING NETWORK DYNAMICS

Combining static regulatory or protein–protein interactions with dynamic data can lead to a better understanding of protein or gene function and can reveal global changes in network topology that hint at higher level cellular organizational principles and functions. The goals of such work often involve finding key regulators of transcription or of dynamic cellular responses in specific conditions. For example, by integrating transcriptional regulatory networks with gene expression data from different conditions, Luscombe *et al.* offer a functional characterization of yeast hub proteins as either ‘transient’ or constitutively active (‘permanent’) [96]. Among the transient hubs, they further distinguish between those responding to programmed changes (such as sporulation or the cell cycle), which tend to involve slower, multi-level response cascades, and those regulating reactive changes to environmental stimuli (such as DNA damage or stress response), which cause rapid activation of many targets.

A different approach to dynamic modeling of transcriptional networks is taken by Chechik *et al.* [97]. Their work focuses on identifying network motifs [98, 99], over-represented substructures (such as feed-forward loops) within the regulatory network that are thought to work together to perform specific functions. The basic approach is not unlike that of some of the pathway-discovery methods described above. Possible ‘timing activity motifs’ (a motif structure and a ‘wiring pattern’ indicating the order of expression onset of the associated elements) are enumerated, and motifs over-represented in a large collection of gene expression time series data sets are identified. Inferred response times are derived from the ‘impulse model’, an idealized model representing a common pattern of gene expression state changes [100]. Analysis of the resulting motifs results in identifying bifurcation points as well as feed-forward and ‘backward activation’ patterns thought to quickly regulate cellular response to changing environmental conditions.

Studies of network motifs can also be informative about the dynamics of signal propagation [101]. In the context of regulatory networks, network motifs provide important clues to how signals might be propagated and regulated [102]. Recently, Ma’ayan *et al.* [103] showed, in the context of hippocampal CA1 neurons, that similar motifs are

also formed as information flows through a signaling network. Interestingly, they observed that ligands causing rapid, transient changes are typically characterized by different motifs than ligands facilitating permanent changes.

The relationship between a position of a gene in a network and its biological properties has been also explored in the context of transcriptional networks. An interesting property of transcriptional networks is their semi-hierarchical structure [104–107]. Therefore, one might ask whether the position of a gene in such structured networks is related to its biological properties. To answer this question, Jothi *et al.* divided the genes in the yeast regulatory network into three levels: ‘top’, ‘core’ and ‘bottom’ [107]. They demonstrated that genes within a network level have similar transcriptional half-lives, abundances and noise levels, but that these properties differ between levels. Elucidating such relationships between network structure and biological properties might provide new insights toward understanding cell dynamics.

Analogous to Luscombe *et al.*’s characterization of transcriptional regulatory networks [96], work by de Lichtenberg *et al.* integrates dynamic data and static protein–protein interaction networks for the purpose of characterizing protein complexes as either transient or constitutively expressed [108]. They find that most complexes consist of both types of proteins, and they suggest that the periodically expressed components control complex activity via a just-in-time assembly mechanism. In contrast, Komurov and White look at related data for condition-dependent and condition-independent (constitutive) expression patterns. They conclude that functional modules (not necessarily complexes) are predominantly comprised either of constitutive or of dynamic proteins, but rarely of both [109]. Differences in the data sources and the types of modules considered are likely to explain the different conclusions of these two papers.

Han *et al.* characterized hubs in the yeast protein–protein interaction network into two groups called ‘party’ and ‘date’ hubs [110]. The expression patterns of party hubs’ interaction partners have a high average correlation with the expression pattern of the hub itself, suggesting that all of the interactions may take place simultaneously or under similar conditions. Date hubs, which have a lower average expression correlation with their neighbors, are thought to interact with different partners under

different conditions. The authors suggest that the date hubs are more likely to be global regulators linking lower-level functional modules comprised of party hubs and their neighbors. Importantly, there are significantly more date hubs in the binary network, whereas party hubs are prevalent in the co-complex networks [111]. There is also some debate about how strongly the party–date analysis depends on the exact expression data sets used [112, 113], suggesting that using additional information to identify static or dynamic hubs may be desirable.

Schmidt and McMahon emphasize that network-related concepts, such as the property of being a hub protein, must be placed in a functional and temporal context [114]. In their study, which focuses on the Clathrin-mediated endocytosis (CME) pathway, they demonstrate that hub connectivity changes over different stages of vesicle formation and suggest that the AP2 hub can only function as a hub when many AP2 molecules form a ‘hub assembly zone’. Arguably, hubs play important roles in network modularity and dynamics [101, 114].

RELATING THE EFFECT OF GENE KNOCKOUTS TO NETWORK TOPOLOGY

Are static, topological properties of a network related to the dynamic behavior of the cell in response to perturbations? Jeong *et al.* observed that high-degree nodes in a protein interaction network tend to correspond to proteins essential for the survival of yeast cells in optimum conditions [115]. This is tested by knocking down the corresponding genes and testing the viability of resulting mutant. Subsequently, the relationship between protein essentiality and global and local topological features of the protein interaction network has been broadly investigated by many authors [116–122]. To answer the question of why such hubs are essential, Jeong and colleagues [115] suggested that over-representation of essential proteins among high-degree nodes can be attributed to the central role hubs play in mediating interactions among numerous, less connected proteins. In contrast, He and Zhang proposed that the majority of proteins are essential due to their involvement in one or more essential protein–protein interactions that are distributed uniformly at random among the network edges [120].

If either of the above two views was confirmed, essential hubs would have played a very special role in network dynamics. However, a recent study of Zotenko *et al.* [122] rejected both of these explanations. Instead, they proposed that the majority of hubs are essential due to their involvement in Essential Complex Biological Modules (ECOBIMs), a group of densely connected proteins with shared biological function that are enriched in essential proteins. This and related studies [123–126] suggest that phenotypic traits such as essentiality have a modular nature. In particular, Zotenko *et al.* showed that, for most data sets and a number of centrality measures, after correcting for correlation between degree and essentiality, essentiality typically correlates with *local* but not *global* centrality measures. Subsequent studies of predicted protein complexes not only confirmed the relationship between essentiality and participation in complexes but also pointed to the relationship between the size of the complex and the essentiality of its members [127]. Interestingly, no relation between essentiality and the number of complexes that include a given gene has been found [126]. However, organizational properties may be related to individual transcript dynamics. While Janga and Babu observed no correlation between essentiality and protein half-life, they did find a correlation between centrality and mRNA half-life [128].

The findings of Zotenko *et al.* explain why only a very weak centrality–lethality relationship [119, 122] was observed in the network resulting from the yeast two-hybrid screen of Ito *et al.* [129]. In addition, a recently constructed, high-quality yeast two-hybrid map showed highly significant clustering between essential proteins [111] but no significant correlation between essentiality and vertex degree. Indeed, one should keep in mind that the yeast two-hybrid assay reports binary interactions only (i.e. direct physical contact as opposed to membership in the same complex), so the size of a complex to which a given protein belongs has relatively low impact on the degree of the corresponding node. Furthermore, large complexes are not always detected by the yeast two-hybrid technique. If the observed relationship between vertex degree and lethality was mostly due to membership in large essential complexes, it is not surprising that the relationship is much weaker in a network comprised only of yeast two-hybrid data.

In contrast to the weak correlation between essentiality and vertex degree, Yu *et al.* [121]

observed that in their large yeast two-hybrid protein interaction network, vertex degree correlates with pleiotropy, the number of phenotypes observed as a consequence of a gene knock-out. Interestingly, such a correlation between the degree and number of phenotypes is not observed in networks constructed from co-purification experiments. These results underscore the importance of considering the experimental technique used to obtain an interaction network when interpreting a network's dynamic properties. Further exploring the relationship between topology and dynamic function, Missiuro *et al.* [116] assess the total information flow through each protein in a network and demonstrate that this information flow score correlates with both essentiality and pleiotropy.

RELATING DOUBLE MUTANT PERTURBATIONS WITH THE PHYSICAL INTERACTOME

Different types of experimental assays may reflect different types of dynamic information. For example, genetic perturbation phenotypes may be inherently condition-specific (e.g. temperature-sensitive lethal mutations), whereas yeast two-hybrid systems create an artificial context to assess the possibility of physical interactions between proteins that may never even appear in proximity to each other *in vivo*. Integrating multiple types of interaction data may help overcome the contextual biases or omissions of particular methods and provide more reliable temporal and contextual information about molecular relationships.

Genetic (or epistatic) interactions occur when the phenotypic impact of one gene is modified by another gene or set of genes. In this review, we will touch upon only symmetric binary interactions, such as synthetic lethals, identified by observing that the composite effect of the simultaneous mutation of two genes has a different phenotypic effect than expected from modeling the combined independent effect of such mutations. Genetic interactions are one means for revealing active information flow in cellular networks. Central methods for uncovering genetic interactions include screens for synthetic lethality and specialized expression Quantitative Trait Loci (eQTL) analysis aimed at identifying genes regulated by several loci. However, one should keep in mind that while synthetic lethal interactions are always between genes, eQTL-based analysis uncovers interacting genetic loci which may

include non-coding regions of the genome. In this section, we survey recent results related to synthetic lethal interactions, and we defer the discussion of interactions between loci to the following section.

A synthetic lethal (or sick) relation between two non-essential genes occurs when simultaneous knockout of the two genes results in cell death (or aggravating genetic impact on cell fitness). Genetic interactions can provide valuable information about protein function and the buffering properties of cellular networks [130]. Considering a hybrid network of genetic and physical interactions provides additional insights into protein function, modular organization of cellular processes and clues as to how these two networks complement each other to achieve the desired robustness to perturbation or response to changing conditions [125, 131–137].

In particular, it has been noted that pairwise interactions between genes often reflect larger relations between whole sets of genes. To capture such dependencies, Kelley and Ideker [131] introduced the *between pathway model* (BPM). This model consists of two sets of proteins that have many genetic interactions *between* the two sets, but few *within* each set, and conversely many physical interactions within each set but few between them. Such a pattern of synthetic-lethal interactions can be explained by a model in which the proteins within each of the two sets function as a pathway (or a protein complex), working together to achieve some goal, and where one set can functionally compensate for the other. Recent work [138] incorporates dynamic information from gene expression data to evaluate the functional coherence and compensation of putative BPMs, focusing attention on the strongest candidates with evident compensatory roles. By integrating temporal data with these topologically derived models, their approach provides additional functional information about the relevant gene sets and the relationships between them.

INTERPERSONAL GENETIC VARIATION AS PERTURBATIONS OF CELLULAR SYSTEMS

Natural genetic variations within and between species can also be viewed as genome-wide ‘perturbations’. These perturbations can occur anywhere in the genomic sequence, not necessarily just within genes. Correlation of genotypic variations with observed phenotypes can provide clues about possible causal relationships between genetic

mutations and the corresponding phenotypic variations. Many recent association results utilize whole genome gene expression data as the phenotype, and thereby such eQTL analysis allows observing the most direct consequences of genetic polymorphism, namely its effects on gene expression. Since the pioneering work of Jansen and Nap [139], eQTL analysis has been applied to a large number of species including yeast [140–143], mouse [143, 144] and human [145–147]. With eQTL analysis, it is possible to elucidate several properties of gene regulation including putative causative relationships between expression variation and genetic loci [148–151], co-regulated gene modules [140, 150, 151–156] and whole regulatory networks [157–161]. Integrative eQTL studies are widely used to identify disease causing mutations [162].

It is important to keep in mind that the large number of gene expression traits and genomic loci poses challenges for both computational efficiency and statistical power. Traditionally, an eQTL study tests the linkage between all genes’ expression and all loci, adding up to millions of individual statistical tests. This leads to unique challenges related to multiple testing adjustment [163]. Recently, Yang *et al.* proposed a graph-theoretical method for performing an eQTL analysis [164] that is able to uncover associations missed by traditional approaches. More work is still needed in developing methods with the statistical power to identify such context-dependent relationships, particularly those that involve multiple genes simultaneously.

To strengthen the signal and to explore additional dependencies between genes and their expression, several of the above-mentioned methods integrate additional information, including protein-protein interaction data, transcription factor-binding sites, co-expression and sequence conservation [154, 155, 161, 165]. Some utilize variants of information flow approaches to uncover pathways connecting putative causal genes and their targets [149,151].

As mentioned above, simultaneous knockdowns of pairs of non-essential genes allow discovery of genetic interactions between genes. Similarly, gene expression variation analyzed in the context of natural genetic variation present in the population is being increasingly used to discover interactions between loci. Epistatic interactions are typically revealed by testing whether the expression of a gene is better explained by a model of genetic variation at one locus or at multiple loci. Epistatic

interactions appear to be persistent among expression trait loci. Brem *et al.* [141] estimated that 16% of heritable transcripts in yeast exhibit epistasis. From a statistical and computational perspective, discovering epistatic interactions on a genome-wide scale is even more challenging than discovering individual eQTLs. Thus, despite significant progress [166], finding epistatic interactions on a genome-wide scale remains a major challenge. Currently, little is known about the dynamics of such interactions. While some work has been done toward identifying gene–environment or gene–context interactions involving individual genes [167], future work relating epistatic genetic relationships with response to exogenous stimuli such as environmental stressors is essential.

CONCLUSIONS

Since the dynamics of protein interactions play a central role in regulating the functioning of cells and organisms, and since high-throughput experimental techniques are increasingly helpful in elucidating interaction dynamics, we expect computational approaches for inferring and analyzing network dynamics to be a cornerstone in future efforts to understand biological systems. Here we have briefly reviewed some recent computational attempts to obtain a dynamic view of the interactome. Many of these approaches have built upon existing static molecular interaction data sets and integrated other types of information to help infer interaction dynamics, both to identify the paths by which information flows through cellular networks and to obtain a more global view of network organizational principles. These other sources have included gene expression data, phenotypic responses to perturbations such as gene knock-outs and information about eQTLs. In the near future, we anticipate that incorporating additional data sources, resulting from advances in mass spectrometry, next-generation sequencing and other high-throughput experimental methods, will help in further elucidating protein interaction and network dynamics. Novel data may include, for example, quantitative protein expression levels, protein localization and modification data, non-coding RNAs and detailed information about epigenetic regulation and the transcriptome. Looking further ahead, it is possible that incorporating future genome-scale data about dynamic responses to changes in DNA structure (e.g. super-coiling and its regulatory effects [168–170]) will

require further expanding our dynamic models of regulatory control [171].

Experimental technologies for uncovering protein interaction dynamics are also being developed and are being applied to increasingly large systems. With respect to detecting protein–protein interactions, promising techniques include fluorescence resonance energy transfer (FRET) [172], which has been used, for example, to map protein interaction in the *Escherichia coli* chemotaxis pathway and determine stimulation induced changes [173]; fluorescent tagging of proteins, followed by rapid isolations and mass spectrometry identification of interactions, which has been used to uncover the dynamics of an alphavirus protein’s host protein interactions [174]; and a protein-fragment complementation assay that has allowed detection of yeast protein interactions within their proper cellular contexts [175]. For regulatory interactions, ChIP-seq technology is likely to be increasingly applied to uncover contextual and temporal variation. As existing technical difficulties with these and other technologies are overcome and larger dynamic networks are elucidated, computational approaches will increasingly switch focus from attempts to infer network dynamics to the analysis of experimentally determined dynamic networks. We expect that the emerging field of dynamic network analysis, which includes algorithmic approaches for modeling and analysis, mostly in the context of social and information networks, will be a good source of analytical methods relevant to dynamic biological networks as well [176–181].

As increasingly diverse high-throughput experimental data sets are gathered, it is likely that novel computational models, or new applications of existing models, will be needed to incorporate the data into predictive quantitative models of the dynamic interactome. While many probabilistic network models have already been applied to incorporate disparate data sets in a single framework, new methods that can intelligently relate different types of interaction data with different sorts of temporal and contextual biases may be helpful. For example, hypergraph models might be used for explicit representations of different types of interaction data. Important questions arising from efforts to merge disparate data types include how to propagate different types of dynamic information among an integrated data set, how to represent explicit bias and how to identify and resolve discrepancies.

An important consideration generally overlooked in the work reviewed here is that temporal variations affecting network dynamics can range over several orders of magnitude. This wide range not only raises questions about increasing and expanding the temporal resolution of experimental methods but also demands the development of appropriate multi-timescale computational methods. Another simplification made by many of the approaches that we have reviewed is that they typically model biological networks with a single node for each protein; in reality, each node represents a population of proteins that may be in different phosphorylation, activation or interaction states. Dynamical systems approaches for analysis and simulation [17, 18] provide a framework for handling these complexities, and integrating these approaches with more high-throughput graph-theoretic methods may be particularly fruitful.

To conclude, we believe that the shift from static to dynamic network analysis is essential for further understanding of molecular systems. Furthermore, we postulate that ignoring dynamic information in future computational network analysis research could turn out to be perilous. Indeed, while the analysis of static protein interaction networks has led to many hypotheses concerning cellular organization, pathways and function, some of these may be challenged by analyses incorporating network dynamics. Thus, we encourage computational biologists who are studying molecular interactions and networks to develop methods that incorporate information about the dynamic nature of cellular systems. After nearly a decade of work focusing on static interactomes, we think it's about time.

Key Points

- Temporally, spatially and contextually dependent molecular interactions play a key role in regulating the functioning of cells and organisms. Incorporating and analyzing these dynamics are the next major challenges for computational interactomics.
- While most large-scale technologies to determine protein interactions do not provide spatial, temporal or contextual information about detected interactions, dynamic views of the interactome can be obtained by integrating static molecular interaction data sets with other types of dynamic data, such as gene expression data, phenotypic responses to perturbations such as gene knock-outs or knock-downs, and information about expression Quantitative Trait Loci (eQTLs).
- It is essential that computational biologists studying molecular networks increasingly shift their focus to develop methods that incorporate information about the dynamic nature of cellular systems.

Acknowledgements

This article was inspired by our involvement in the Pacific Symposium on Biocomputing session 'Dynamics of Biological Networks', which we are co-organizing with Tanya Berger-Wolf (University of Illinois). We thank her for many illuminating discussions, and the organizers of the PSB conference for supporting our session. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

FUNDING

Intramural Research Program of the National Institutes of Health and National Library of Medicine (to T.M.P.); National Science Foundation grant CCF-0542187, National Institutes of Health grant GM076275 and National Institutes of Health Center of Excellence grant P50 GM071508 (M.S.); National Institutes of Health grants LM009411 and HD058880 (to D.K.S.).

References

1. Ito T, Chiba T, Ozawa R, *et al.* A comprehensive two-hybrid analysis to explore the yeast protein interactome. *Proc Natl Acad Sci USA* 2001;**98**:4569–74.
2. Uetz P, Giot L, Cagney G, *et al.* A comprehensive analysis of protein–protein interactions in *Saccharomyces cerevisiae*. *Nature* 2000;**403**:623–7.
3. Ho Y, Gruhler A, Heilbut A, *et al.* Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature* 2002;**415**:180–3.
4. Gavin AC, Aloy P, Grandi P, *et al.* Proteome survey reveals modularity of the yeast cell machinery. *Nature* 2006;**440**:631–6.
5. Krogan NJ, Cagney G, Yu H, *et al.* Global landscape of protein complexes in the yeast *Saccharomyces cerevisiae*. *Nature* 2006;**440**:637–43.
6. Giot L, Bader JS, Brouwer C, *et al.* A protein interaction map of *Drosophila melanogaster*. *Science* 2003;**302**:1727–36.
7. Li S, Armstrong CM, Bertin N, *et al.* A map of the interactome network of the metazoan *C. elegans*. *Science* 2004;**303**:540–3.
8. Rual JF, Venkatesan K, Hao T, *et al.* Towards a proteome-scale map of the human protein–protein interaction network. *Nature* 2005;**437**:1173–8.
9. Rain JC, Selig L, De Reuse H, *et al.* The protein–protein interaction map of *Helicobacter pylori*. *Nature* 2001;**409**:211–5.
10. Harbison CT, Gordon DB, Lee TI, *et al.* Transcriptional regulatory code of a eukaryotic genome. *Nature* 2004;**431**:99–104.
11. Lee TI, Rinaldi NJ, Robert F, *et al.* Transcriptional regulatory networks in *Saccharomyces cerevisiae*. *Science* 2002;**298**:799–804.
12. Ptacek J, Devgan G, Michaud G, *et al.* Global analysis of protein phosphorylation in yeast. *Nature* 2005;**438**:679–84.

13. Aittokallio T, Schwikowski B. Graph-based methods for analysing networks in cell biology. *Brief Bioinform* 2006;**7**: 243–55.
14. Zhu X, Gerstein M, Snyder M. Getting connected: analysis and principles of biological networks. *Genes Dev* 2007;**21**: 1010–24.
15. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* 2000;**28**:27–30.
16. Tyson JJ, Chen K, Novak B. Network dynamics and cell physiology. *Nat Rev Mol Cell Biol* 2001;**2**:908–16.
17. Tyson JJ, Chen KC, Novak B. Sniffers, buzzers, toggles and blinkers: dynamics of regulatory and signaling pathways in the cell. *Curr Opin Cell Biol* 2003;**15**:221–31.
18. de Jong H. Modeling and simulation of genetic regulatory systems: a literature review. *J Comput Biol* 2002;**9**:67–103.
19. Ruths D, Muller M, Tseng JT, *et al.* The signaling petri net-based simulator: a non-parametric strategy for characterizing the dynamics of cell-specific signaling networks. *PLoS Comput Biol* 2008;**4**:e1000005.
20. Zielinski R, Przytycki PF, Zheng J, *et al.* The crosstalk between EGF, IGF, and Insulin cell signaling pathways—computational and experimental analysis. *BMC Syst Biol* 2009;**3**:88.
21. Rubinstein A, Gurevich V, Kasulin-Boneh Z, *et al.* Faithful modeling of transient expression and its application to elucidating negative feedback regulation. *Proc Natl Acad Sci USA* 2007;**104**:6241–6.
22. Aldridge BB, Saez-Rodriguez J, Muhlich JL, *et al.* Fuzzy logic analysis of kinase pathway crosstalk in TNF/EGF/insulin-induced signaling. *PLoS Comput Biol* 2009;**5**: e1000340.
23. Thakar J, Piloni M, Kirimanjeswara G, *et al.* Modeling systems-level regulation of host immune responses. *PLoS Comput Biol* 2007;**3**:e109.
24. De Jong H, Gouze JL, Hernandez C, *et al.* Qualitative simulation of genetic regulatory networks using piecewise-linear models. *Bull Math Biol* 2004;**66**:301–40.
25. Albert I, Thakar J, Li S, *et al.* Boolean network simulations for life scientists. *Source Code Biol Med* 2008;**3**:16.
26. Zhang R, Shah MV, Yang J, *et al.* Network model of survival signaling in large granular lymphocyte leukemia. *Proc Natl Acad Sci USA* 2008;**105**:16308–13.
27. Fields S, Song O. A novel genetic system to detect protein–protein interactions. *Nature* 1989;**340**:245–6.
28. Collins MO, Choudhary JS. Mapping multiprotein complexes by affinity purification and mass spectrometry. *Curr Opin Biotechnol* 2008;**19**:324–30.
29. Kim TH, Ren B. Genome-wide analysis of protein–DNA interactions. *Annu Rev Genomics Hum Genet* 2006;**7**:81–102.
30. Johnson DS, Mortazavi A, Myers RM, *et al.* Genome-wide mapping of *in vivo* protein–DNA interactions. *Science* 2007;**316**:1497–1502.
31. Ni L, Bruce C, Hart C, *et al.* Dynamic and complex transcription factor binding during an inducible response in yeast. *Genes Dev* 2009;**23**:1351–63.
32. Jakobsen JS, Braun M, Astorga J, *et al.* Temporal ChIP-on-chip reveals Biniou as a universal regulator of the visceral muscle transcriptional network. *Genes Dev* 2007;**21**: 2448–60.
33. Steffen M, Petti A, Aach J, *et al.* Automated modelling of signal transduction networks. *BMC Bioinform* 2002;**3**:34.
34. Scott J, Ideker T, Karp RM, *et al.* Efficient algorithms for detecting signaling pathways in protein interaction networks. *J Comput Biol* 2006;**13**:133–44.
35. Supper J, Spangenberg L, Planatscher H, *et al.* BowTieBuilder: modeling signal transduction pathways. *BMC Syst Biol* 2009;**3**:67.
36. Csete M, Doyle J. Bow ties, metabolism and disease. *Trends Biotechnol* 2004;**22**:446–50.
37. Banks E, Nabieva E, Peterson R, *et al.* NetGrep: fast network schema searches in interactomes. *Genome Biol* 2008;**9**: R138.
38. Banks E, Nabieva E, Chazelle B, *et al.* Organization of physical interactomes as uncovered by network schemas. *PLoS Comput Biol* 2008;**4**:e1000203.
39. Zotenko E, Guimaraes KS, Jothi R, *et al.* Decomposition of overlapping protein complexes: a graph theoretical method for analyzing static and dynamic protein associations. *Algorithms Mol Biol* 2006;**1**:7.
40. Gavril F. The intersection graphs of subtrees in trees are exactly the chordal graphs. *J Comb Theory: Ser B* 1974;**16**: 47–56.
41. Monma CL, Wei VK. Intersection graphs of paths in a tree. *J Comb Theory: Ser B* 1986;**41**:141–81.
42. Kelley BP, Sharan R, Karp RM, *et al.* Conserved pathways within bacteria and yeast as revealed by global protein network alignment. *Proc Natl Acad Sci USA* 2003;**100**: 11394–9.
43. Koyuturk M, Kim Y, Topkara U, *et al.* Pairwise alignment of protein interaction networks. *J Comput Biol* 2006;**13**: 182–99.
44. Sharan R, Suthram S, Kelley RM, *et al.* Conserved patterns of protein interaction in multiple species. *Proc Natl Acad Sci USA* 2005;**102**:1974–9.
45. Flannick J, Novak A, Srinivasan BS, *et al.* Graemlin: general and robust alignment of multiple large interaction networks. *Genome Res* 2006;**16**:1169–81.
46. Singh R, Xu J, Berger B. Global alignment of multiple protein interaction networks with application to functional orthology detection. *Proc Natl Acad Sci USA* 2008;**105**: 12763–8.
47. Sharan R, Ideker T. Modeling cellular machinery through biological network comparison. *Nat Biotechnol* 2006;**24**: 427–33.
48. Ideker T, Ozier O, Schwikowski B, *et al.* Discovering regulatory and signalling circuits in molecular interaction networks. *Bioinformatics* 2002;**18**(Suppl 1):S233–40.
49. Cabusora L, Sutton E, Fulmer A, *et al.* Differential network expression during drug and stress response. *Bioinformatics* 2005;**21**:2898–2905.
50. Guo Z, Li Y, Gong X, *et al.* Edge-based scoring and searching method for identifying condition-responsive protein–protein interaction sub-network. *Bioinformatics* 2007;**23**: 2121–8.
51. Ulitsky I, Shamir R. Identification of functional modules using network topology and high-throughput data. *BMC Syst Biol* 2007;**1**:8.
52. Yeang CH, Ideker T, Jaakkola T. Physical network models. *J Comput Biol* 2004;**11**:243–62.
53. Yeang CH, Mak HC, McCuine S, *et al.* Validation and refinement of gene-regulatory pathways on a network of physical interactions. *Genome Biol* 2005;**6**:R62.

54. Ourfali O, Shlomi T, Ideker T, *et al.* SPINE: a framework for signaling-regulatory pathway inference from cause-effect experiments. *Bioinformatics* 2007;**23**:i359–66.
55. Singh R, Berger B. Influence flow: integrating pathway-specific RNAi data and protein interaction data. PLoS track at ISMB/ECCB. Cambridge, UK: PLoS, 2007.
56. Tu Z, Argmann C, Wong KK, *et al.* Integrating siRNA and protein-protein interaction data to identify an expanded insulin signaling network. *Genome Res* 2009;**19**:1057–67.
57. Jones TR, Carpenter AE, Lamprecht MR, *et al.* Scoring diverse cellular morphologies in image-based screens with iterative feedback and machine learning. *Proc Natl Acad Sci USA* 2009;**106**:1826–31.
58. Moffat J, Grueneberg DA, Yang X, *et al.* A lentiviral RNAi library for human and mouse genes applied to an arrayed viral high-content screen. *Cell* 2006;**124**:1283–98.
59. Beyer A, Bandyopadhyay S, Ideker T. Integrating physical and genetic maps: from genomes to interaction networks. *Nat Rev Genet* 2007;**8**:699–710.
60. Bumgarner RE, Yeung KY. Methods for the inference of biological pathways and networks. *Methods Mol Biol* 2009;**541**:225–45.
61. Friedman N. Inferring cellular networks using probabilistic graphical models. *Science* 2004;**303**:799–805.
62. D’Haeseleer P, Liang S, Somogyi R. Genetic network inference: from co-expression clustering to reverse engineering. *Bioinformatics* 2000;**16**:707–26.
63. Pe’er D, Regev A, Elidan G, *et al.* Inferring subnetworks from perturbed expression profiles. *Bioinformatics* 2001;**17**(Suppl 1):S215–24.
64. Margolin AA, Nemenman I, Basso K, *et al.* ARACNE: an algorithm for the reconstruction of gene regulatory networks in a mammalian cellular context. *BMC Bioinform* 2006;**7**(Suppl 1):S7.
65. Margolin AA, Wang K, Lim WK, *et al.* Reverse engineering cellular networks. *Nat Protoc* 2006;**1**:662–71.
66. Friedman N, Linial M, Nachman I, *et al.* Using Bayesian networks to analyze expression data. *J Comput Biol* 2000;**7**:601–20.
67. Hartemink AJ, Gifford DK, Jaakkola TS, *et al.* Using graphical models and genomic expression data to statistically validate models of genetic regulatory networks. *Pac Symp Biocomput* 2001;**6**:422–33.
68. Sachs K, Perez O, Pe’er D, *et al.* Causal protein-signaling networks derived from multiparameter single-cell data. *Science* 2005;**308**:523–9.
69. Baym M, Bakal C, Perrimon N, *et al.* High resolution modeling of cellular signaling networks. *RECOMB* 2008. Singapore: ;257–71.
70. Friedman N, Murphy K, Russell S. Learning the structure of dynamic probabilistic networks. *Proceedings of the 14th Conference on Uncertainty in Artificial Intelligence*. San Mateo: Morgan Kaufmann, 1998;139–47.
71. Murphy K, Mian S. Modeling Gene Expression Data using Bayesian Networks. *Technical Report, Computer Science Division*. Berkeley, CA: University of California.
72. Ong IM, Glasner JD, Page D. Modelling regulatory pathways in *E. coli* from time series expression profiles. *Bioinformatics* 2002;**18**(Suppl 1):S241–8.
73. Kim SY, Imoto S, Miyano S. Inferring gene networks from time series microarray data using dynamic Bayesian networks. *Brief Bioinform* 2003;**4**:228–35.
74. Butte AJ, Kohane IS. Mutual information relevance networks: functional genomic clustering using pairwise entropy measurements. *Pac Symp Biocomput* 2000;**5**:415–26.
75. Lee HK, Hsu AK, Sajdak J, *et al.* Coexpression analysis of human genes across many microarray data sets. *Genome Res* 2004;**14**:1085–94.
76. Stuart JM, Segal E, Koller D, *et al.* A gene-coexpression network for global discovery of conserved genetic modules. *Science* 2003;**302**:249–55.
77. Choi JK, Yu U, Yoo OJ, *et al.* Differential coexpression analysis using microarray data and its application to human cancer. *Bioinformatics* 2005;**21**:4348–55.
78. Kostka D, Spang R. Finding disease specific alterations in the co-expression of genes. *Bioinformatics* 2004;**20**(Suppl 1):i194–9.
79. Fang G, Kuang R, Pandey G, *et al.* Subspace differential coexpression analysis: problem definition and a general approach. *Pac Sympos Biocomput* 2010;**15**:145–56.
80. Ahmed A, Xing EP. Recovering time-varying networks of dependencies in social and biological studies. *Proc Natl Acad Sci USA* 2009;**106**:11878–83.
81. Song L, Kolar M, Xing EP. KELLER: estimating time-varying interactions between genes. *Bioinformatics* 2009;**25**:i128–36.
82. Cheng Y, Church GM. Biclustering of expression data. *Proc Int Conf Intell Syst Mol Biol* 2000;**8**:93–103.
83. Madeira SC, Oliveira AL. Biclustering algorithms for biological data analysis: a survey. *IEEE/ACM Trans Comput Biol Bioinform* 2004;**1**:24–45.
84. Segal E, Shapira M, Regev A, *et al.* Module networks: identifying regulatory modules and their condition-specific regulators from gene expression data. *Nat Genet* 2003;**34**:166–76.
85. Bonneau R, Reiss DJ, Shannon P, *et al.* The Inferelator: an algorithm for learning parsimonious regulatory networks from systems-biology data sets *de novo*. *Genome Biol* 2006;**7**:R36.
86. Lemmens K, De Bie T, Dhollander T, *et al.* DISTILLER: a data integration framework to reveal condition dependency of complex regulons in *Escherichia coli*. *Genome Biol* 2009;**10**:R27.
87. de Bivort B, Huang S, Bar-Yam Y. Empirical multiscale networks of cellular regulation. *PLoS Comput Biol* 2007;**3**:1968–78.
88. Cooke EJ, Savage RS, Wild DL. Computational approaches to the integration of gene expression: ChIP-chip and sequence data in the inference of gene regulatory networks. *Semin Cell Dev Biol* 2009;**20**:863–8.
89. Bromberg KD, Ma’ayan A, Neves SR, *et al.* Design logic of a cannabinoid receptor signaling network that triggers neurite outgrowth. *Science* 2008;**320**:903–9.
90. Baugh LR, Hill AA, Claggett JM, *et al.* The homeodomain protein PAL-1 specifies a lineage-specific regulatory network in the *C. elegans* embryo. *Development* 2005;**132**:1843–54.
91. Liao JC, Boscolo R, Yang YL, *et al.* Network component analysis: reconstruction of regulatory signals in biological systems. *Proc Natl Acad Sci USA* 2003;**100**:15522–7.

92. Seok J, Xiao W, Moldawer LL, *et al.* A dynamic network of transcription in LPS-treated human subjects. *BMC Syst Biol* 2009;**3**:78.
93. Ye C, Galbraith SJ, Liao JC, *et al.* Using network component analysis to dissect regulatory networks mediated by transcription factors in yeast. *PLoS Comput Biol* 2009;**5**: e1000311.
94. Ernst J, Vainas O, Harbison CT, *et al.* Reconstructing dynamic regulatory maps. *Mol Syst Biol* 2007;**3**:74.
95. Ucar D, Beyer A, Parthasarathy S, *et al.* Predicting functionality of protein-DNA interactions by integrating diverse evidence. *Bioinformatics* 2009;**25**:i137–44.
96. Luscombe NM, Babu MM, Yu H, *et al.* Genomic analysis of regulatory network dynamics reveals large topological changes. *Nature* 2004;**431**:308–12.
97. Chechik G, Oh E, Rando O, *et al.* Activity motifs reveal principles of timing in transcriptional control of the yeast metabolic network. *Nat Biotechnol* 2008;**26**:1251–9.
98. Shen-Orr SS, Milo R, Mangan S, *et al.* Network motifs in the transcriptional regulation network of *Escherichia coli*. *Nat Genet* 2002;**31**:64–8.
99. Milo R, Itzkovitz S, Kashtan N, *et al.* Superfamilies of evolved and designed networks. *Science* 2004;**303**: 1538–42.
100. Chechik G, Koller D. Timing of gene expression responses to environmental changes. *J Comput Biol* 2009;**16**: 279–90.
101. Alexander RP, Kim PM, Emonet T, *et al.* Understanding modularity in molecular networks requires dynamics. *Sci Signal* 2009;**2**:pe44.
102. Alon U. Network motifs: theory and experimental approaches. *Nat Rev Genet* 2007;**8**:450–61.
103. Ma'ayan A, Jenkins SL, Neves S, *et al.* Formation of Regulatory Patterns During Signal Propagation in a Mammalian Cellular Network. *Science* 2005;**309**:1078–83.
104. Balazsi G, Barabasi AL, Oltvai ZN. Topological units of environmental signal processing in the transcriptional regulatory network of *Escherichia coli*. *Proc Natl Acad Sci USA* 2005;**102**:7841–6.
105. Ma HW, Buer J, Zeng AP. Hierarchical structure and modules in the *Escherichia coli* transcriptional regulatory network revealed by a new top-down approach. *BMC Bioinform* 2004;**5**:199.
106. Yu H, Gerstein M. Genomic analysis of the hierarchical structure of regulatory networks. *Proc Natl Acad Sci USA* 2006;**103**:14724–31.
107. Jothi R, Balaji S, Wuster A, *et al.* Genomic analysis reveals a tight link between transcription factor dynamics and regulatory network architecture. *Mol Syst Biol* 2009;**5**:294.
108. de Lichtenberg U, Jensen LJ, Brunak S, *et al.* Dynamic complex formation during the yeast cell cycle. *Science* 2005;**307**:724–27.
109. Komurov K, White M. Revealing static and dynamic modular architecture of the eukaryotic protein interaction network. *Mol Syst Biol* 2007;**3**:110.
110. Han JD, Bertin N, Hao T, *et al.* Evidence for dynamically organized modularity in the yeast protein-protein interaction network. *Nature* 2004;**430**:88–93.
111. Yu H, Braun P, Yildirim MA, *et al.* High-quality binary protein interaction map of the yeast interactome network. *Science* 2008;**322**:104–10.
112. Batada NN, Reguly T, Breitkreutz A, *et al.* Stratus not altocumulus: a new view of the yeast protein interaction network. *PLoS Biol* 2006;**4**:e317.
113. Batada NN, Reguly T, Breitkreutz A, *et al.* Still stratus not altocumulus: further evidence against the date/party hub distinction. *PLoS Biol* 2007;**5**:e154.
114. Schmid EM, McMahon HT. Integrating molecular and network biology to decode endocytosis. *Nature* 2007;**448**: 883–8.
115. Jeong H, Mason SP, Barabasi AL, *et al.* Lethality and centrality in protein networks. *Nature* 2001;**411**:41–2.
116. Missiuro PV, Liu K, Zou L, *et al.* Information flow analysis of interactome networks. *PLoS Comput Biol* 2009;**5**: e1000350.
117. Yu H, Greenbaum D, Xin Lu H, *et al.* Genomic analysis of essentiality within protein networks. *Trends Genet* 2004;**20**: 227–31.
118. Hahn MW, Kern AD. Comparative genomics of centrality and essentiality in three eukaryotic protein-interaction networks. *Mol Biol Evol* 2005;**22**:803–6.
119. Batada NN, Hurst LD, Tyers M. Evolutionary and physiological importance of hub proteins. *PLoS Comput Biol* 2006;**2**:e88.
120. He X, Zhang J. Why do hubs tend to be essential in protein networks? *PLoS Genet* 2006;**2**:e88.
121. Yu H, Kim PM, Sprecher E, *et al.* The importance of bottlenecks in protein networks: correlation with gene essentiality and expression dynamics. *PLoS Comput Biol* 2007;**3**:e59.
122. Zotenko E, Mestre J, O'Leary DP, *et al.* Why do hubs in the yeast protein interaction network tend to be essential: reexamining the connection between the network topology and essentiality. *PLoS Comput Biol* 2008;**4**:e1000140.
123. Fraser HB, Plotkin JB. Using protein complexes to predict phenotypic effects of gene mutation. *Genome Biol* 2007;**8**: R252.
124. Hart GT, Lee I, Marcotte ER. A high-accuracy consensus map of yeast protein complexes reveals modular nature of gene essentiality. *BMC Bioinform* 2007;**8**:236.
125. Bandyopadhyay S, Kelley R, Krogan NJ, *et al.* Functional maps of protein complexes from quantitative genetic interaction data. *PLoS Comput Biol* 2008;**4**:e1000065.
126. Pache RA, Babu MM, Aloy P. Exploiting gene deletion fitness effects in yeast to understand the modular architecture of protein complexes under different growth conditions. *BMC Syst Biol* 2009;**3**:74.
127. Wang H, Kakaradov B, Collins SR, *et al.* A complex-based reconstruction of the *S. cerevisiae* interactome. *Mol Cell Proteomics* 2009;**8**:1361–81.
128. Janga SC, Babu MM. Transcript stability in the protein interaction network of *Escherichia coli*. *Mol Biosyst* 2009;**5**: 154–62.
129. Ito T, Chiba T, Ozawa R, *et al.* A comprehensive two-hybrid analysis to explore the yeast protein interactome. *Proc Natl Acad Sci USA* 2001;**98**:4569–74.
130. Tong AH, Evangelista M, Parsons AB, *et al.* Systematic genetic analysis with ordered arrays of yeast deletion mutants. *Science* 2001;**294**:2364–8.
131. Kelley R, Ideker T. Systematic interpretation of genetic interactions using protein networks. *Nat Biotechnol* 2005;**23**:561–6.

132. Collins SR, Miller KM, Maas NL, *et al.* Functional dissection of protein complexes involved in yeast chromosome biology using a genetic interaction map. *Nature* 2007;**446**:806–10.
133. Roguev A, Wiren M, Weissman JS, *et al.* High-throughput genetic interaction mapping in the fission yeast *Schizosaccharomyces pombe*. *Nat Methods* 2007;**4**:861–6.
134. Ulitsky I, Shamir R. Pathway redundancy and protein essentiality revealed in the *Saccharomyces cerevisiae* interaction networks. *Mol Syst Biol* 2007;**3**:104.
135. Ma X, Tarone AM, Li W. Mapping genetically compensatory pathways from synthetic lethal interactions in yeast. *PLoS ONE* 2008;**3**:e1922.
136. Ulitsky I, Shlomi T, Kupiec M, *et al.* From E-MAPs to module maps: dissecting quantitative genetic interactions using physical interactions. *Mol Syst Biol* 2008;**4**:209.
137. Brady A, Maxwell K, Daniels N, *et al.* Fault tolerance in protein interaction networks: stable bipartite subgraphs and redundant pathways. *PLoS ONE* 2009;**4**:e5364.
138. Hescott BJ, Leiserson MDM, Cowen LJ, *et al.* Evaluating between-pathway models with expression data. *Res Comp Mol Biol Proc* 2009;**5541**:372–85.
139. Jansen RC, Nap JP. Genetical genomics: the added value from segregation. *Trends Genet* 2001;**17**:388–91.
140. Yvert G, Brem RB, Whittle J, *et al.* Trans-acting regulatory variation in *Saccharomyces cerevisiae* and the role of transcription factors. *Nat Genet* 2003;**35**:57–64.
141. Brem RB, Kruglyak L. The landscape of genetic complexity across 5,700 gene expression traits in yeast. *Proc Natl Acad Sci U S A* 2005;**102**:1572–7.
142. Brem RB, Storey JD, Whittle J, *et al.* Genetic interactions between polymorphisms that affect gene expression in yeast. *Nature* 2005;**436**:701–3.
143. Klose J, Nock C, Herrmann M, *et al.* Genetic analysis of the mouse brain proteome. *Nat Genet* 2002;**30**:385–93.
144. Chesler EJ, Lu L, Shou S, *et al.* Complex trait analysis of gene expression uncovers polygenic and pleiotropic networks that modulate nervous system function. *Nat Genet* 2005;**37**:233–42.
145. Monks SA, Leonardson A, Zhu H, *et al.* Genetic inheritance of gene expression in human cell lines. *Am J Hum Genet* 2004;**75**:1094–105.
146. Cheung VG, Spielman RS, Ewens KG, *et al.* Mapping determinants of human gene expression by regional and genome-wide association. *Nature* 2005;**437**:1365–9.
147. Stranger BE, Forrest MS, Clark AG, *et al.* Genome-wide associations of gene expression variation in humans. *PLoS Genet* 2005;**1**:e78.
148. Kulp DC, Jagalur M. Causal inference of regulator–target pairs by gene mapping of expression phenotypes. *BMC Genomics* 2006;**7**:125.
149. Tu Z, Wang L, Arbeitman MN, *et al.* An integrative approach for causal gene identification and gene regulatory pathway inference. *Bioinformatics* 2006;**22**:e489–96.
150. Sun W, Yu T, Li KC. Detection of eQTL modules mediated by activity levels of transcription factors. *Bioinformatics* 2007;**23**:2290–7.
151. Suthram S, Beyer A, Karp RM, *et al.* eQED: an efficient method for interpreting eQTL associations using protein networks. *Mol Syst Biol* 2008;**4**:162.
152. Li H, Chen H, Bao L, *et al.* Integrative genetic analysis of transcription modules: towards filling the gap between genetic loci and inherited traits. *Hum Mol Genet* 2006;**15**:481–92.
153. Sampson JN, Self SG. Identifying trait clusters by linkage profiles: application in genetical genomics. *Bioinformatics* 2008;**24**:958–64.
154. Zhu J, Zhang B, Smith EN, *et al.* Integrating large-scale functional genomic data to dissect the complexity of yeast regulatory networks. *Nat Genet* 2008;**40**:854–61.
155. Lee SI, Dudley AM, Drubin D, *et al.* Learning *a priori* on regulatory potential from eQTL data. *PLoS Genet* 2009;**5**:e1000358.
156. Litvin O, Causton HC, Chen BJ, *et al.* Modularity and interactions in the genetics of gene expression. *Proc Natl Acad Sci USA* 2009;**106**:6441–6.
157. Zhu J, Lum PY, Lamb J, *et al.* An integrative genomics approach to the reconstruction of gene networks in segregating populations. *Cytogenet Genome Res* 2004;**105**:363–74.
158. Bing N, Hoeschele I. Genetical genomics analysis of a yeast segregant population for transcription network inference. *Genetics* 2005;**170**:533–42.
159. Li H, Lu L, Manly KF, *et al.* Inferring gene transcriptional modulatory relations: a genetical genomics approach. *Hum Mol Genet* 2005;**14**:1119–25.
160. Liu B, de la Fuente A, Hoeschele I. Gene network inference via structural equation modeling in genetical genomics experiments. *Genetics* 2008;**178**:1763–76.
161. Huang Y, Zheng J, Przytycka T. Discovery of regulatory mechanisms by genome-wide from gene expression variation by eQTL analysis. In: Lonardi S, Chen JY, (eds). *Biological Data Mining*. Boca Raton, FL: CRC Press, 2009; 205–28.
162. Schadt EE. Exploiting naturally occurring DNA variation and molecular profiling data to dissect disease and drug response traits. *Curr Opin Biotechnol* 2005;**16**:647–54.
163. Storey JD, Tibshirani R. Statistical significance for genomewide studies. *Proc Natl Acad Sci USA* 2003;**100**:9440–5.
164. Huang Y, Wuchty S, Ferdig MT, *et al.* Graph theoretical approach to study eQTL: a case study of *Plasmodium falciparum*. *Bioinformatics* 2009;**25**:i15–20.
165. Schadt EE, Lamb J, Yang X, *et al.* An integrative genomics approach to infer causal associations between gene expression and disease. *Nat Genet* 2005;**37**:710–7.
166. Cordell HJ. Genome-wide association studies: detecting gene–gene interactions that underlie human diseases. *Nat Rev Genet* 2009;**10**:392–404.
167. Hunter DJ. Gene–environment interactions in human diseases. *Nat Rev Genet* 2005;**6**:287–98.
168. Kouzine F, Levens D. Supercoil-driven DNA structures regulate genetic transactions. *Front Biosci* 2007;**12**:4409–23.
169. Kouzine F, Sanford S, Elisha-Feil Z, *et al.* The functional response of upstream DNA to dynamic supercoiling *in vivo*. *Nat Struct Mol Biol* 2008;**15**:146–54.
170. Cheung KJ, Badarinarayana V, Selinger DW, *et al.* A microarray-based antibiotic screen identifies a regulatory role for supercoiling in the osmotic stress response of *Escherichia coli*. *Genome Res* 2003;**13**:206–15.
171. Marr C, Geertz M, Hutt MT, *et al.* Dissecting the logical types of network control in gene expression profiles. *BMC Syst Biol* 2008;**2**:18.

172. Piston DW, Kremers GJ. Fluorescent protein FRET: the good, the bad and the ugly. *Trends Biochem Sci* 2007;**32**: 407–14.
173. Kentner D, Sourjik V. Dynamic map of protein interactions in the *Escherichia coli* chemotaxis pathway. *Mol Syst Biol* 2009;**5**:238.
174. Cristea IM, Carroll JW, Rout MP, et al. Tracking and elucidating alphavirus-host protein interactions. *J Biol Chem* 2006;**281**:30269–78.
175. Tarassov K, Messier V, Landry CR, et al. An *in vivo* map of the yeast protein interactome. *Science* 2008;**320**:1465–70.
176. Kleinberg J. Small-world phenomena and the dynamics of information. *Adv Neural Inform Process Syst (NIPS)* 2001;**14**: 431–8.
177. Breiger RL, Carley KM, Pattison P. *Dynamic Social Network Analysis: Workshop Summary and Papers*. Washington, DC: The National Academies Press, 2002.
178. Kempe D, Kleinberg J, Kumar A. Connectivity and inference problems for temporal networks. *J Comput Syst Sci* 2002;**64**:820–42.
179. Kempe D, Kleinberg J, Tardos E. Maximizing the spread of influence through a social network. *Proceeding of the 9th ACM SIGKDD Intl. Conf. on Knowledge Discovery and Data Mining*. New York, NY: ACM Press, 2003; 137–46.
180. Leskovec J, Kleinberg J, Faloutsos C. Graphs over time: densification laws, shrinking diameters and possible explanations. *Proceedings of the 11th ACM SIGKDD International Conference on Knowledge Discovery in Data Mining*. IL USA: ACM, Chicago, 2005;177–87.
181. Tantipathananandh C, Berger-Wolf T, Kempe D. A framework for community identification in dynamic social networks. *Proceedings of the 13th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*. CA, USA: ACM, San Jose, 2007;717–26.